

Product Datasheet

[Anti-Glypican 4 \(GPC4\) \[IPI-mGPC4.9\]](#)

[Applications](#) | [Antibody Details](#) | [Antigen Details](#) | [References](#)

Overview

Antigen	Glypican 4 (GPC4)
Immunogen	Purified recombinant fragment of Mouse Glypican 4 (GPC4), corresponding to AA: 23-528.
Host/isotype	Rabbit/IgG1
Clonality	Recombinant monoclonal
Clone name	IPI-mGPC4.9
RRID	AB_3101863
IPI ID	TAB0010818-013-002
Specificity	GPC4, GPC6 (weak); Does not recognize other GPCs
Species reactivity	Mouse and human
Amount	100 µg
Concentration	1 mg/mL
Purification	Expressed in ExpiCHO cells and affinity purified using Protein A
Storage buffer	PBS, pH 7.4 with 0.02% ProClin 300
Shipping	Shipped on blue ice at +4C
Storage	Store at +4C for up to 3 months. For long-term storage, aliquot and store at -20C. Avoid multiple freeze/thaw cycles.

IPI Tested Applications[‡]

Application	Tested concentration	Result	Reference
Cell Display	5 µg/mL	Positive	https://doi.org/10.57733/addgene.nscs9i
Flow	5 µg/mL	Positive	https://doi.org/10.57733/addgene.y1j609
IF	1 µg/mL	Positive	https://doi.org/10.57733/addgene.45gbjl

[‡] Not suitable for WB application.

Community Data*


Application	Lab	Reference
IHC	James Trimmer, Ph.D., UC Davis	https://doi.org/10.57733/addgene.0vnll9

* Supporting Data is generated by external partner labs, in the process of evaluating IPI antibody panels.

FOR RESEARCH USE ONLY - NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Applications


Cell Display



Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] recognizes mouse and human GPC4. Histogram from FACS analysis on ExpiCHO cells transfected with mouse or human glypican 4 (red), or WY0041 negative control (gray). Cells expressing mouse (left panel) or human (right panel) GPC4 were labeled with Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] (Addgene #222516) and Alexa Fluor 647 F(ab')₂ Fragment Goat Anti-Rabbit IgG Fc (Jackson ImmunoResearch, 111-606-046). Labeled cells were analyzed with an Intellicyt iQue Screener Plus flow cytometer. Histograms were generated and normalized to mode using FlowJo™ v10.10. doi:

<https://doi.org/10.57733/addgene.nscs9j>


Flow cytometry



Histogram overlays from FACS analysis on ExpiCHO cells transfected with human and mouse glypicans (GPC1-GPC6). Cells expressing GPCs were labeled with Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] (Addgene #222516) and Alexa Fluor 647 F(ab')₂ Fragment Goat Anti-Rabbit IgG Fc (Jackson ImmunoResearch, 111-606-046). Labeled cells were analyzed with an Intellicyt iQue Screener Plus flow cytometer. Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] is specific for human and mouse GPC4 (violet lines with fill) and is weakly cross-reactive with mouse GPC6 (blue line with fill). Histograms were normalized to mode and displayed as an overlay with half offset using FlowJo™ v10.10. Bottom table summarizes GPC cross-reactivity. Abbreviations: glypican (GPC), human (Hu), mouse (Mo), strong (++), weak (+), none (.). doi: <https://doi.org/10.57733/addgene.y1j609>

FOR RESEARCH USE ONLY - NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Immunofluorescence



Immunofluorescence (IF) of ExpiCHO cells transfected with human GPC4. ExpiCHO cells were transiently transfected with GFP (transfection control) and human GPC4. 100,000 cells were fixed to a 96-well glass bottom plate using 4% PFA at 37C for 15 min. Cells were permeabilized with 0.1% Triton X-100 at RT for 10 min and then blocked with 5% BSA, 5% serum, and 0.01% Triton X-100 at room temperature for 30 min. Cells were incubated with 1 μ g/mL Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] (Addgene #222516). Alexa Fluor 647 F(ab')₂ fragment goat anti-rabbit IgG, Fc fragment-specific (Jackson ImmunoResearch, 111-606-046) was used as the secondary antibody. Nuclear staining was performed using 0.5 mg/mL DAPI. Cells were imaged for GFP (green), DAPI (blue), and GPC4 (red) using an EVOS M7000 microscope at 10x magnification. doi: <https://doi.org/10.57733/addgene.45gbjl>

Antibody Details

Antibody design and production

Human variable domains for the heavy and light chain of the FAB fragment used in yeast display were grafted onto the constant CH1, CH2 and CH3 domains of rabbit IgG. The chimera human/rabbit IgG1 construct was recombinantly expressed in ExpiCHO cells, using pTipi2.1 as the expression vector. The antibody was purified by affinity chromatography using protein A (XYZ) and acid elution, followed by immediate buffer exchange using 1 x PBS buffer pH 7.4.

Sequence information

Heavy chain and light chain amino acid sequences are available upon request after purchase. [Contact us](#) to request.

Antibody Characterization

LC-MS: Intact mass analysis via LC-MS methods allows for confirmation antibody mass, and to identify any product-related variants such as glycosylation. Before conducting intact mass analysis via LC-MS, the antibody was reduced to its heavy chain (HC) and light chain (LC). This process allows for confirmation of the masses corresponding to the amino acid sequences of both chains.


	HC MW (Da) <i>Calculated</i>	HC MW (Da) <i>Observed</i>	HC MW (Da) <i>Delta</i>	LC MW (Da) <i>Calculated</i>	LC MW (Da) <i>Observed</i>	LC MW (Da) <i>Delta</i>
IPI-mGPC4.9	50080.15	50085.40	5.25	22952.44	22952.57	0.13

Heavy Chain (HC) Mass Calculation: The calculated molecular weight (MW) of the HC is derived by adding the mass of the unmodified HC amino acid sequence to the mass of the predominant N-glycan form (G0F), which is 1444.5 Da. This calculation assumes that the intrachain disulfide bonds remain intact. For HCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), resulting in a deduction of 17.03 Da from the total mass. Additionally, for HCs with a C-terminal lysine (K), the mass of K (128.09 Da) is also subtracted.

Light Chain (LC) Mass Calculation: The calculated molecular weight (MW) of the LC is obtained from the mass of the unmodified LC amino acid sequence, assuming that the intrachain disulfide bonds are not reduced. For LCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), leading to a deduction of 17.03 Da from the total mass. For LCs with a C-terminal lysine (K), the mass of K (128.09 Da) is subtracted as well.

FOR RESEARCH USE ONLY - NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Size Exclusion Chromatography (SEC): SEC is a protein purification technique that separates molecules based on size.



High-performance liquid chromatography-size exclusion chromatography (HPLC-SEC) profile. 5 μ g of antibody (1 mg/mL) was analyzed. 1x PBS, pH 7.4 was used as the mobile phase. Protein standards ranging from 15-600 kDa were used for calibration (Sigma, 69385) and Lox1.9 antibody was run as an internal control. Resulting peaks were analyzed using Agilent OpenLab CDS software to determine total peak area and percentage purity.

	Purity	Result
IPI-mGPC4.9	$\geq 95\%$	Pass

FOR RESEARCH USE ONLY - NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Antigen Details

Immunogen design:

cDNA of Mouse Glypican 4 (GPC4) with C-terminal His-, Avi-, and HA-tags was produced in transiently transfected Expi293F cells and purified from culture supernatant by Ni-NTA affinity purification followed by size-exclusion chromatography.

Immunogen sequences:

Mouse GPC4 (AA: 23-528)

AELKSKSCSEVRRRLVSKGFNKNDAPLYEINGDHLKICPQDYTCSSQEMEEKYSLQSKDDFKTVVSEQC
NHLQAIFASRYKKFDEFFKELLENAEKSLNDMFVKTYGHLYMQNSELFKDLFVELKRYVAGNVNLEEML
NDFWARLLERMFRVLNSQYHFTDEYLECVSKYTEQLKPFQGDVPRKLLQVTRAFVAARTFAQGLAVARD
VSKVSVWNPTAQCTHALLKMIYCSHCRGLVTVKPCYNYCSNIMRGCLANQGDLDFEWNNFIDAMLMV
AERLEGPFNIESVMDPIDVKISDAIMNMQDNSVQVSQKVFQGCQPPKPLPAGRISRSISESAFSARFRPY
HPEQRPTTAAGTSLDRLVTDVKEKLLQAKKFWSSLPSTVCNDERMAAGNENEDDCWNGKKGKSRYLFA
VTGNLANQGNNPEVQVDTSKPDILILRQIMALRVMTSKMKNAYNGNDVDFDISDESSGEGSGSGCEY
QQCPSEFEYNATDHSGKSANEKADSGSGHHHHHHHHHHHSGGGLNDIFEAQKIEWHEGSGDYKDDDD
K

Sequence information:

HUGO: [MGI:104902](#)
Uniprot: [P51655](#)
Refseq: [NM_008150.2](#)

Structural information:

Topology: Glycosylphosphatidylinositol (GPI) Anchored
PDB IDs: None
AlphaFold: [AF-P51655-F1](#)

Expression profiles:

Human Protein Atlas [ENSG00000076716-GPC4](#)

References

1. Z. Anderson, H. Li, T. Riedel, H Zhu, R. Meijers, and D. Moshinsky. (2024). Flow Cytometry for Anti-Glypican 4 (GPC4) [IPI-GPC4.9]. Addgene. <https://doi.org/10.57733/addgene.nscs9i>
2. Z. Anderson, H. Li, T. Riedel, H Zhu, R. Meijers, and D. Moshinsky. (2024). Flow Cytometry for Anti-Glypican 4 (GPC4) [IPI-GPC4.9]. Addgene. <https://doi.org/10.57733/addgene.y1j609>
3. H. Li, Z. Anderson, T. Riedel, and D. Moshinsky. (2024). IF for Anti-Glypican 4 (GPC4) [IPI-mGPC4.9]. Addgene. <https://doi.org/10.57733/addgene.45gbjl>
4. J. Trimmer. (2024). IHC for Anti-Glypican 4 (GPC4) [IPI-mGPC4.9]. Addgene. <https://doi.org/10.57733/addgene.0vnll9>

How to cite this antibody:

Anti-Glypican 4 (GPC4) [IPI-mGPC4.9] - from Institute for Protein Innovation (IPI) (Addgene #222516; <http://n2t.net/addgene:222516>; RRID: AB_3101863).

If you publish research with this product, please [let us know](#) so that we can cite your paper.