

Product Datasheet

[Anti-Glypican 2 \(GPC2\) \[IPI-mGPC2.31\]](#)

[Applications](#) | [Antibody Details](#) | [Antigen Details](#) | [References](#)

Overview

Antigen	Glypican 2 (GPC2)
Immunogen	Purified recombinant fragment of Mouse Glypican 2 (GPC2), corresponding to AA: 24-578.
Host/isotype	Rabbit/IgG
Clonality	Recombinant monoclonal
Clone name	IPI-mGPC2.31
RRID	AB_3697522
IPI ID	TAB0015525-013-002
Specificity	GPC2; Does not recognize other GPCs
Species reactivity	Mouse
Amount	100 µg
Concentration	1 mg/mL
Purification	Expressed in HEK293 cells and affinity purified using Protein A
Storage buffer	PBS, pH 7.4
Shipping	Shipped on blue ice at +4C
Storage	Store at +4C for up to 3 months. For long-term storage, aliquot and store at -20C. Avoid multiple freeze/thaw cycles.

IPI Tested Applications[‡]

Application	Tested concentration	Result	Reference
Flow	0.66-100 µg/mL	Positive	https://doi.org/10.57733/addgene.p1tc82
IF – Binding	1 µg/mL	Positive	https://doi.org/10.57733/addgene.hmvk5r
IF – Specificity	1 µg/mL	Positive	https://doi.org/10.57733/addgene.w9uv3e

[‡] Not suitable for WB application.

Community Data*


Application	Lab	Reference
IHC	James Trimmer, Ph.D., UC Davis/NeuroMab	https://doi.org/10.57733/addgene.rz7aaf

* Supporting Data is generated by external partner labs, in the process of evaluating IPI antibody panels.

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Applications

Flow cytometry




Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] (Addgene #240998) is specific for mouse GPC2 in flow cytometry. Histogram from FACS analysis on ExpiCHO cells transfected with mouse or human GPC2 (red), or B7H3 negative control (gray). Cells expressing mouse (left panel) or human (right panel) GPC2 were labeled with Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] and Alexa Fluor 647 F(ab')₂ goat anti-rabbit IgG Fc fragment (Jackson ImmunoResearch, 111-606-046). Labeled cells were analyzed with an Intellicyrt iQue Screener Plus flow cytometer. Histograms were generated and normalized to mode using FlowJo™ v10.10. **Note:** A reduced mouse GPC2 right shift was observed and caused by low cell expression of the antigen. doi: <https://doi.org/10.57733/addgene.p1tc82>

EC₅₀ (data not shown): A fourteen-point titration of antibody concentrations, ranging from 660 nM (0.1 mg/mL) to 4.42 pM with a 1:2.5 dilution factor was performed against human and mouse GPC2. Results showed specificity towards mouse GPC2 only, with an observed EC₅₀ value of 0.68 nM for mouse GPC2.

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE


Immunofluorescence (IF) – Species Reactivity



Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] (Addgene #240998) is specific for GPC2 in immunofluorescence. A) Immunofluorescence (IF) of ExpiCHO cells transfected with human and mouse GPC2. Human GPC2 was co-transfected with GFP (human transfection control) and mouse GPC2 was co-transfected with BFP (mouse transfection control). Confocal images taken at 40X magnification on the ImageXpress confocal HT.ai microscope. Cells were imaged for GFP (green), BFP (blue), and GPC2 (magenta). B) Combined quantification of multiple images of the same transfected cells taken at 10X magnification. GFP- or BFP- positive cells were identified using a custom module in the IN Carta image analysis software, then the mean fluorescence intensity (MFI) of the far-red channel for each cell, representing IPI-mGPC2.31 staining, was recorded. Each histogram displays the number of cells with MFIs ranging from below 450 (background fluorescence) to 20000. IPI-mGPC2.31 staining of human (left) and mouse (right) GPC2 are shown in the top row, and compared to a positive (left) and negative (right) control in the bottom row. For both panels, IPI-mGPC2.31 was used at 1 μ g/mL (1:1,000 dilution). doi: <https://doi.org/10.57733/addgene.hmvk5r>

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Immunofluorescence (IF) – Target Specificity



Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] (Addgene #240998) is specific for mouse GPC2 in immunofluorescence. (Top) Immunofluorescence (IF) of ExpiCHO cells transfected with human and mouse GPC1-GPC6. Human GPC targets were co-transfected with GFP (human transfection control) and mouse GPC targets were co-transfected with BFP (mouse transfection control). Widefield images were taken at 10X magnification on an imageXpress confocal HT.ai microscope. (Bottom) Each graph depicts the combined quantification of multiple images of the same transfected cells taken at 10X magnification. GFP- or BFP-positive cells were identified using a custom module in the InCarta software, then the mean fluorescence intensity (MFI) of the far-red channel for each cell, representing IPI-mGPC2.31 staining, was recorded. Each histogram displays the number of cells with MFIs ranging from below 450 (background fluorescence) to 20000. IPI-mGPC2.31 staining of human and mouse variants of each GPC family member is compared on the top and bottom rows. For all panels, IPI-mGPC2.31 was used at 1 µg/mL (1:1,000 dilution). doi: <https://doi.org/10.57733/addgene.w9uv3e>

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Antibody Details

Antibody design and production

Human variable domains for the heavy and light chain of the FAB fragment used in yeast display were grafted onto the constant CH1, CH2 and CH3 domains of rabbit IgG. The chimera human/rabbit IgG1 construct was recombinantly expressed in Expi HEK293 cells, using pTipi2.1 as the expression vector. The antibody was purified by affinity chromatography using protein A (XYZ) and acid elution, followed by immediate buffer exchange using 1 x PBS buffer pH 7.4.

Sequence information

Heavy chain and light chain amino acid sequences are available upon request after purchase. [Contact us](#) to request.

Antibody Characterization

LC-MS: Intact mass analysis via LC-MS methods allows for confirmation antibody mass, and to identify any product-related variants such as glycosylation. Before conducting intact mass analysis via LC-MS, the antibody was reduced to its heavy chain (HC) and light chain (LC). This process allows for confirmation of the masses corresponding to the amino acid sequences of both chains.


	HC MW (Da) <i>Calculated</i>	HC MW (Da) <i>Observed</i>	HC MW (Da) <i>Delta</i>	LC MW (Da) <i>Calculated</i>	LC MW (Da) <i>Observed</i>	LC MW (Da) <i>Delta</i>
IPI-mGPC2.31	49758.60	49762.10	3.50	22952.44	22952.19	-0.25

Heavy Chain (HC) Mass Calculation: The calculated molecular weight (MW) of the HC is derived by adding the mass of the unmodified HC amino acid sequence to the mass of the predominant N-glycan form (G0F), which is 1444.5 Da. This calculation assumes that the intrachain disulfide bonds remain intact. For HCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), resulting in a deduction of 17.03 Da from the total mass. Additionally, for HCs with a C-terminal lysine (K), the mass of K (128.09 Da) is also subtracted.

Light Chain (LC) Mass Calculation: The calculated molecular weight (MW) of the LC is obtained from the mass of the unmodified LC amino acid sequence, assuming that the intrachain disulfide bonds are not reduced. For LCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), leading to a deduction of 17.03 Da from the total mass. For LCs with a C-terminal lysine (K), the mass of K (128.09 Da) is subtracted as well.

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE


Size Exclusion Chromatography (SEC): SEC is a protein purification technique that separates molecules based on size.



	RT (min)	Width (min)	Area	Height	Area %	Result
IPI-mGPC2.31	2.966	0.1819	611.0452	55.9848	90.5915	Pass

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Surface Plasmon Resonance (SPR)



	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	K_D (M)	Res. sd
IPI-mGPC2.31	2.0×10^5	1.1×10^{-3}	5.8×10^{-9}	37.8

Surface Plasmon Resonance (SPR) kinetics analysis of the interaction between Anti-GPC2 [IPI-mGPC2.31] and mouse GPC2. SPR binding kinetics were measured on a Carterra LSA using HC30M chips (Carterra, cat. #4279) at 25 °C. Goat anti-rabbit IgG Fc (Jackson ImmunoResearch, cat. #111-005-046) was immobilized via amine coupling, and test antibodies were captured using a 96-channel print-head. Antigens (400 nM to five lower concentrations, 2-fold dilutions) were injected in antigen buffer (20 mM HEPES pH 7.4, 150 mM NaCl, 1 mM CaCl₂, 1 mM MgCl₂, 0.005% Tween 80) with 300 s association/dissociation phases and acid regeneration. Data (reference/buffer subtracted, smoothed) were globally fit to a 1:1 Langmuir model to derive k_a , k_d , and K_D using Carterra Kinetics software v1.9.2.44.63, and replotted in OriginPro 2023b. Results show a high-affinity and specific binding event between the antibody and antigen.

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Antigen Details

Immunogen design:

cDNA of Mouse Glypican 2 (GPC2) with C-terminal Avi-tag was produced in transiently transfected Expi293F cells and purified from culture supernatant by Ni-NTA affinity purification followed by size-exclusion chromatography.

Immunogen sequences:

>Mouse GPC2 (AA: 24-578):

SEAKVVRSCAETRQVLGARGYSLNLIPPSLISGEHLQVCPQEYTCCSSETEQKLIRDAEVTFRGLVEDSGS
FLIHTLAARHRKFNEFFREMLISQHSQAQLFSHSYGRLYSQHAVIFNSLFSGLRDYYEKSGEGLDDTLAD
FWAQLLERAFPLHPQYSFPPDFLLCLTRLTSTADGSLQPFQDSPRRRLRLQISRALVAARALVQGLETGR
NVVSEALKVPVLEGCRQALMRLIGCPLCRGVPSLMPCRGFCLNVAHGCLSSRGLEPEWGGYLDGLLLL
AEKLQGPFSFELAAESIGVKISEGLMHLQENSVKVSQKVFQECGTPHPVQSRSRVAPAPREEASRSWVA
SAEEERPTTAAGTNLHRLVWELRERLSRVRFWAGLPVTVCGDSRMAADLSQETAPCWTGVGRGRYM
SPVVVGSLSNEQLHNPDLTSSPDVPTRRVRLHLRAATARMKAAALGQDLDMHDADEDASGSGGGQQY
ADDWKAGAVPVPPARPPRPVPPRRDGLGVRGGSGSARYNQGRSRNGGGGGSGGGGSPAPPELLGGP
SVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
LAQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVNTLCLVKGFYPSDIA
VEWESNGQPENNYKTTTPVLDSDGSFFLNSTLTVDKSRWQQGNVFCSSVMHEALANHYTQKLSLSLSPG
KSGSGLNDIFEAQKIEWHE

Sequence information:

HUGO: MGI:1919201
Uniprot: Q8BKV1
Refseq: NM_172412.3

Structural information:

Topology: Glycosylphosphatidylinositol (GPI) Anchored
PDB IDs: 6WJL;7T62
AlphaFold: AF-Q8BKV1-F1

Expression profiles:

Human Protein Atlas ENSMUSG00000029510

FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

References

1. J. Trimmer. (2025). Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] in Immunohistochemistry (Rat). Addgene.
<https://doi.org/10.57733/addgene.rz7aaf>

How to cite this antibody:

Anti-Glypican 2 (GPC2) [IPI-mGPC2.31] - from Institute for Protein Innovation (IPI) (Addgene #240998; <http://n2t.net/addgene:240998>; RRID: AB_3697522).

If you publish research with this product, please [let us know](#) so that we can cite your paper.