Product Datasheet

Applications | Antibody Details | Antigen Details | References

Overview

Antigen	ROBO2
Immunogen	Purified recombinant fragment of Human ROBO2, corresponding to AA: 22-860.
Host/isotype	Rabbit/IgG
Clonality	Recombinant monoclonal
Clone name	IPI-ROBO2.47
RRID	AB_3720933
IPI ID	TAB0010617-013
Specificity	ROBO2; Weak cross-reactivity with ROBO3.
Species	Human and mouse
Concentration	1 mg/mL
Purification	Expressed in HEK293T cells and affinity purified using Protein A
Storage Buffer	PBS, pH 7.4
Shipping	Shipped on blue ice at +4 °C
Storage	Store at +4C for up to 3 months. For long-term storage, aliquot and store at -20C. Avoid multiple freeze/thaw cycles.

IPI Tested Applications ‡

Applications	Tested Concentration	Result	Reference
Flow Cytometry	0.66-100 µg/mL	Positive	https://doi.org/10.57733/addgene.hiazwc
IF - Binding	1 µg/mL	Positive	https://doi.org/10.57733/addgene.gibssr
IF - Specificity	1 µg/mL	Positive	https://doi.org/10.57733/addgene.8j600d

‡ Not suitable for WB application.

Community Data *

Application	Lab	Reference
ICC/IF	Homaira Nawabi, Ph.D., University Grenoble-Alpes	https://doi.org/10.57733/addgene.b9zkoq

* Community data is generated by external partner labs, in the process of evaluating IPI antibody panels.

FOR RESEARCH USE ONLY – NOT INTENDED FOR DIAGNOSTIC or THERAPEUTIC USE

Applications

Cell Display



Anti-ROBO2 [IPI-ROBO2.47] (Addgene #251719) recognizes human and mouse ROBO2.

Histogram from FACS analysis on ExpiCHO cells transfected with human or mouse ROBO2 (red), or B7H3 negative control (gray). Cells expressing human (left panel) or mouse (right panel) ROBO2 were labeled with Anti-ROBO2 [IPI-ROBO2.47] and Alexa Fluor 647 F(ab')2 goat anti-rabbit IgG Fc fragment (Jackson ImmunoResearch, 111-606-046). Labeled cells were analyzed with an Intellicyt iQue Screener Plus flow cytometer. Histograms were generated and normalized to mode using FlowJo™ v10.10.

doi: https://doi.org/10.57733/addgene.hiazwc

Immunofluorescence (IF) – Binding



Anti-ROBO2 [IPI-ROBO2.47] (Addgene#251719) shows binding to human and mouse ROBO2.

A) Immunofluorescence (IF) of ExpiCHO cells transfected with human (Green/GFP) and mouse (Blue/BFP) ROBO2 stained with IPI-ROBO2.47 (magenta). Confocal images taken at 40X magnification on the ImageXpress confocal HT.ai microscope. Scale bar = 20 uM. B) Combined quantification of multiple images of the same transfected cells taken at 10X magnification. GFP- or BFP-positive cells were identified via the neural network CellPose, then the mean fluorescence intensity (MFI) of the far-red channel for each cell, representing IPI-ROBO2.47 staining, was recorded. Each histogram displays the number of cells with MFIs ranging from below 100 (background fluorescence) to 22000. IPI-ROBO2.47 staining of human and mouse ROBO2 is shown in the top row, and compared to a positive (left) and negative (right) control in the bottom row. For both panels, IPI-ROBO2.47 was used at 1 ug/mL (1:1,000 dilution).

doi: https://doi.org/10.57733/addgene.gibssr

Immunofluorescence (IF) - Target Specificity



Family Crossreactivity

	ROBO1		ROBO2		ROBO3		ROBO4	
	Hu	Mo	Hu	Mo	Hu	Mo	Hu	Mo
IPI-ROBO2.47			++	++	+	+		

Strong ++
Weak +
None

Anti-ROBO2 [IPI-ROBO2.47] (Addgene#251719) shows significant binding only to human and mouse ROBO2.

Each graph depicts the combined quantification of multiple images of the same transfected cells taken at 10X magnification. GFP or BFP-positive cells were identified via the neural network CellPose, then the mean fluorescence intensity (MFI) of the far-red channel for each cell, representing IPI-ROBO2.47 staining, was recorded. Each histogram displays the number of cells with MFIs ranging from below 100 (background fluorescence) to 22000. IPI-ROBO2.47 staining of human and mouse variants of each ROBO family member is compared on the top and bottom rows. To test family-wide cross-reactivity, IPI-ROBO2.47 was used at 1 ug/mL (1:1000 dilution).

doi: https://doi.org/10.57733/addgene.8j600d

Antibody Details

Antibody Design and Production

Human variable domains for the heavy and light chain of the FAB fragment used in yeast display were grafted onto the constant CH1, CH2 and CH3 domains of rabbit IgG. The chimera human/rabbit IgG1 construct was recombinantly expressed in HEK293 cells, using pTipi2.1 as the expression vector. The antibody was purified by affinity chromatography using protein A (XYZ) and acid elution, followed by immediate buffer exchange using 1 x PBS buffer pH 7.4.

Sequence information: Heavy chain and light chain amino acid sequences are available upon request after purchase. Contact us to request.

Antibody Characterization

LC-MS

LC-MS: Intact mass analysis via LC-MS methods allows for confirmation antibody mass, and to identify any product-related variants such as glycosylation. Before conducting intact mass analysis via LC-MS, the antibody was reduced to its heavy chain (HC) and light chain (LC). This process allows for confirmation of the masses corresponding to the amino acid sequences of both chains.

LC-MS Results

LC Calculated	LC Observed	LC Delta	HC Calculated	HC Observed	HC Delta
22618.94	22618.54	-0.40	49424.74	49431.63	6.89

Heavy Chain (HC) Mass Calculation: The calculated molecular weight (MW) of the HC is derived by adding the mass of the unmodified HC amino acid sequence to the mass of the predominant N-glycan form (G0F), which is 1444.5 Da. This calculation assumes that the intrachain disulfide bonds remain intact. For HCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), resulting in a deduction of 17.03 Da from the total mass. Additionally, for HCs with a Cterminal lysine (K), the mass of K (128.09 Da) is also subtracted.

Light Chain (LC) Mass Calculation: The calculated molecular weight (MW) of the LC is obtained from the mass of the unmodified LC amino acid sequence, assuming that the intrachain disulfide bonds are not reduced. For LCs with an N-terminal glutamine (Q), the mass of Q is converted to pyroglutamic acid (PyroGlu), leading to a deduction of 17.03 Da from the total mass. For LCs with a C-terminal lysine (K), the mass of K (128.09 Da) is subtracted as well.

Size Exclusion Chromatography (SEC)

5 µL of 1±0.2 mg/mL of sample was analyzed with TSKgel® SuperSW mAb HTP (4 µm) HPLC Column in 1X PBS, pH 7.4 as the mobile phase.



Clone	RT (min)	Width (min)	Area	Height	Area %	Result
IPI-ROBO2.47	2.90	0.24	141.27	8.53	100.00	Pass

Interpretation: SEC-QC pass. Main peak with greater than 90% purity and within retention time of 2.6 to 3.26 mins.

Antigen Details

Immunogen sequence

>HumanROBO2(AA:22-860):SRLRQEDFPPRIVEHPSDVIVSKGEPTTLNCKAEGRPTPTIEWYKDGERVETDKDDPRSH
RMLLPSGSLFFLRIVHGRRSKPDEGSYVCVARNYLGEAVSRNASLEVALLRDDFRQNPTDVVVAAGEPAILECQPPRGH
PEPTIYWKKDKVRIDDKEERISIRGGKLMISNTRKSDAGMYTCVGTNMVGERDSDPAELTVFERPTFLRRPINQVVLEEEA
VEFRCQVQGDPQPTVRWKKDDADLPRGRYDIKDDYTLRIKKTMSTDEGTYMCIAENRVGKMEASATLTVRAPPQFVVRP
RDQIVAQGRTVTFPCETKGNPQPAVFWQKEGSQNLLFPNQPQQPNSRCSVSPTGDLTITNIQRSDAGYYICQALTVAGSI
LAKAQLEVTDVLTDRPPPIILQGPANQTLAVDGTALLKCKATGDPLPVISWLKEGFTFPGRDPRATIQEQGTLQIKNLRISD
TGTYTCVATSSSGETSWSAVLDVTESGATISKNYDLSDLPGPPSKPQVTDVTKNSVTLSWQPGTPGTLPASAYIIEAFSQS
VSNSWQTVANHVKTTLYTVRGLRPNTIYLFMVRAINPQGLSDPSPMSDPVRTQDISPPAQGVDHRQVQKELGDVLVRLH
NPVVLTPTTVQVTWTVDRQPQFIQGYRVMYRQTSGLQATSSWQNLDAKVPTERSAVLVNLKKGVTYEIKVRPYFNEFQG
MDSESKTVRTTEEAPSAPPQSVTVLTVGSYNSTSISVSWDPPPPDHQNGIIQEYKIWCLGNETRFHINKTVDAAIRSVIIGG
LFPGIQYRVEVAASTSAGVGVKSEPQPIIIGRRNEVVITENNNSITEQITDVVKQPGHHHHHHHHHHHGSGGLNDIFEAQKIE
WHEGSGYPYDVPDYA

Sequence information:

HUGO:	10250
Uniprot:	Q9HCK4
Refseq:	XM_047448669.1

Structural information:

Topology:	Single-pass type I membrane protein
PDB IDs:	1UEM; 1UJT; 2EDJ; 5NOI; 6I9S; 6IAA
Alphafold:	AF-Q9HCK4-F1

Expression profile:

Human Protein Atlas:	ENSG00000185008

References

1. Z. Anderson, H. Li, T. Riedel, H. Zhu and D. Moshinsky. (2026). Flow Cytometry for Anti-ROBO2 [IPI-ROBO2.47]. Addgene. https://doi.org/10.57733/addgene.hiazwc
2. A. Morano, T. Riedel, and D. Moshinsky. (2026). ICC/IF for Anti-ROBO2 [IPI-ROBO2.47] in binding assay. Addgene. https://doi.org/10.57733/addgene.gibssr
3. A. Morano, T. Riedel, and D. Moshinsky. (2026). ICC/IF for Anti-ROBO2 [IPI-ROBO2.47] in specificity assay. Addgene. https://doi.org/10.57733/addgene.8j600d
4. A. Saintpierre and H. Nawabi. (2026). ICC/IF for Anti-ROBO2 [IPI-ROBO2.47]. Addgene. https://doi.org/10.57733/addgene.b9zkoq

How to cite this antibody:

Anti-ROBO2 [IPI-ROBO2.47] - from Institute for Protein Innovation (IPI) (Addgene #251719; xttp://n2t.net/addgene:251719; RRID: AB_3720933).

If you publish research with this product, please let us know so that we can cite your paper.